

Nucleon matrix elements and baryon masses in the Dirac orbital model

Yu.A.Simonov* and M.A.Trusov†

Institute of Theoretical and Experimental Physics, Moscow, Russia

Abstract

Using the expansion of the baryon wave function in a series of products of single quark bispinors (Dirac orbitals), the nonsinglet axial and tensor charges of a nucleon are calculated. The leading term yields $g_A = 1.27$ in good agreement with experiment. Calculation is essentially parameter-free and depends only on the strong coupling constant value α_s . The importance of lower Dirac bispinor component, yielding 18% to the wave function normalization is stressed. As a check, the baryon decuplet masses in the formalism of this model are also computed using standard values of the string tension σ and the strange quark mass m_s ; the results being in a good agreement with experiment.

*e-mail: simonov@itep.ru

†e-mail: trusov@itep.ru

Axial and tensor charges of a nucleon are important to characterize the basic structure of the nucleon as composed of strongly coupled quarks [1]–[4]. Whereas many baryon characteristics can be reasonably obtained in relativized quark models where all relativistic effects are treated via Salpeter equation [5] and spin corrections, the axial and tensor charges are sensitive to the Dirac structure of quark wave functions, and in particular, as will be shown below, to the negative energy components. The most systematic treatment of relativistic baryon system can be done in the three-body Bethe–Salpeter formalism, however in this approach a rigorous derivation ends up in the system of more than 20 integral equations, and therefore, a drastic simplification is needed to make realistic calculations. Recently this kind of approach was developed in [6], where quark-quark interactions have been properly parametrised. However the relativistic objects like negative energy component admixture η (see below) are very sensitive to the form of interactions, and in [6] η has come out rather small. In our paper we choose another and much simpler approach, which allows to treat all Dirac components properly including lower ones, and to work out the resulting η (and axial and tensor charges) in a transparent way.

First of all, let us recall some formulas of the spin relativistic theory. Consider a polarized free fermion with spin $\frac{1}{2}$. The fermion polarization could be described by a spacelike pseudovector a^μ , which reduces in the fermion rest frame to the unit 3-vector along which the fermion spin projection is equal to $+1/2$. The pseudovector a^μ obviously meets conditions: $a \cdot a = -1$, $a \cdot p = 0$, where p is the fermion 4-momentum. The corresponding Michel–Wighimann fermion density matrix is

$$\rho = \frac{\gamma p + m}{4m} [1 - \gamma^5(\gamma a)] \quad (1)$$

where m is the fermion mass. Note, that here and below we propose the plane wave amplitude being normalized to unity: $\bar{u}u = 1$. Using the density matrix (1) the pseudovector a^μ could be also defined in an arbitrary frame as the averaged value of the Pauli–Lubansky operator:

$$\langle \hat{W}^\mu \rangle = \frac{1}{2}a^\mu, \quad \hat{W}^\mu = -\frac{1}{2m}\varepsilon^{\mu\nu\rho\sigma}\hat{J}_{\nu\rho}\hat{P}_\sigma, \quad \varepsilon^{0123} = +1. \quad (2)$$

where $\hat{J}_{\mu\nu}$ is the angular momentum operator.

Axial and tensor current operators are defined as follows:

$$\hat{j}_\mu^A = \hat{\psi}\gamma_\mu\gamma^5\hat{\psi}, \quad \hat{j}_{\mu\nu}^T = \hat{\psi}\sigma_{\mu\nu}\gamma^5\hat{\psi} \quad (3)$$

where $\sigma_\mu = \frac{1}{2}[\gamma_\mu, \gamma_\nu]$ (both currents are hermitean). It is easy to check¹ that for a plane wave

$$\langle \hat{j}_\mu^A \rangle = \bar{u}\gamma_\mu\gamma^5u = -a_\mu, \quad \langle \hat{j}_{\mu\nu}^T \rangle = \bar{u}\sigma_{\mu\nu}\gamma^5u = \frac{1}{m}(a^\mu p^\nu - a^\nu p^\mu) \quad (4)$$

Now, consider a polarized nucleon, e.g. proton, with momentum p and polarization a , and define two quantities — the axial charge g_A and the tensor charge g_T in terms of average values of quark axial and tensor currents similarly to (4), as follows:

$$\langle p, a | \hat{u}\gamma_\mu\gamma^5\hat{u} - \hat{d}\gamma_\mu\gamma^5\hat{d} | p, a \rangle = -g_A a_\mu, \quad (5)$$

¹The simplest way to derive these relations is to work in the fermion rest frame and to choose the Oz axis direction being parallel to the space part of the pseudovector a

$$\left\langle p, a \left| \hat{u} \sigma_{\mu\nu} \gamma^5 \hat{u} - \hat{d} \sigma_{\mu\nu} \gamma^5 \hat{d} \right| p, a \right\rangle = \frac{g_T}{m} (a^\mu p^\nu - a^\nu p^\mu) \quad (6)$$

where m is the proton mass.

In the limit of exact isotopic symmetry (which we assume further), the isotopic structure of the axial current is $2\hat{I}_3$, and its matrix element must coincide with that one of the current from the same isotopic multiplet with structure I_+ taken between proton and neutron. The latter transition arises in neutron β -decay and we conclude that the constant g_A must coincide with the neutron decay axial constant G_A , which is very well known experimentally² $G_A = 1.27$. For the tensor charge, unfortunately, we have no well established experimental value, although it could be in principle measured as the first momentum of the proton transversity distribution function $h_1(x)$ [7].

As for the theoretical calculations of the charges, it is well known that the nonrelativistic quark model predicts $g_A = \frac{5}{3}$, while for massless relativistic quarks in the MIT bag model, one obtains much smaller values $g_A = 1.09$, both in disagreement with experiment. As for the tensor charge g_T , various theoretical estimations give disparate results ranging from 0.89 to 1.45, see [8] for Refs. and discussions. All this certainly stimulates further study of the question.

The approach proposed in this paper could be applied not only for analysis of nucleon properties, but also for the calculation of constants of hyperon semileptonic decays, which values are quite important for the accurate determination of the CKM matrix elements (see, e.g., [9]). We plan to develop a systematic analysis of baryon properties in the framework of Dirac orbital expansion in future papers; the first part of which is reported below.

To construct the baryon wave function, one starts with the Hamiltonian [10] obtained in the instantaneous approximation from the general Bethe-Salpeter equation:

$$\hat{H}\Psi(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) = E\Psi, \quad \hat{H} = \sum_{i=1}^3 \hat{H}_i + \Delta H \quad (7)$$

with

$$\hat{H}_i = \mathbf{p}_{(i)} \boldsymbol{\alpha}_{(i)} + \beta_{(i)} (m_i + M(\mathbf{r}_i)) \quad (8)$$

where $M(\mathbf{r}_i)$ in the limit of vanishing gluon correlation length is

$$M = \sigma |\mathbf{r}_i - \mathbf{r}_X| \quad (9)$$

σ is the string tension, and $\mathbf{r}_X$ is the string-junction coordinate. Here ΔH contains perturbative gluon exchanges. We expand the baryon wave function in a series of products of quark eigenfunctions $\psi_n^{(i)} = \begin{pmatrix} v^{(i)} \\ w^{(i)} \end{pmatrix}$, namely [11]

$$\Psi(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) = \sum_{\{n_i\}} \prod_{i=1}^3 \psi_{n_i}^{(i)}(\mathbf{r}_i) C_{n_1 n_2 n_3} \quad (10)$$

In what follows we shall consider the leading valence approximation for the nucleon keeping only the first term in (10), $\Psi \rightarrow \Psi_0$, which contains the ground state S -wave

²Our choice of the G_A sign corresponds to [1, 2]

Dirac orbitals $|u_\lambda\rangle$ and $|d_\lambda\rangle$ for u and d quarks with spins up and down. So, for the proton polarized along the axis Oz , one has:

$$|p \uparrow\rangle = \sqrt{\frac{1}{18}} [-2(|u \uparrow u \uparrow d \downarrow\rangle + \text{perm.}) + (|u \uparrow u \downarrow d \uparrow\rangle + \text{perm.})] \quad (11)$$

This expression has the same form as in the standard $SU(4)$ or $SU(6)$ model [3] except for the bispinor contents of $|u_\lambda\rangle$ and $|d_\lambda\rangle$. In (11) u_λ and d_λ orbitals depends on space coordinates via the only function χ_λ in accordance with the isotopic symmetry principle. Note, the proton state (11) with S -wave orbitals evidently has the average 3-momentum being equal to zero:

$$\langle \mathbf{P} \rangle = \sum \langle \mathbf{p}_i \rangle = \mathbf{0} \quad (12)$$

where $\mathbf{p}_i$ is the 3-momentum of the i^{th} quark.

Insertion (11) into (5) and (6) yields³:

$$g_A = \langle p \uparrow | \hat{u}^+ \Sigma_3 \hat{u} - \hat{d}^+ \Sigma_3 \hat{d} | p \uparrow \rangle = +\frac{4}{3} \langle \chi_\uparrow | \Sigma_3 | \chi_\uparrow \rangle - \frac{1}{3} \langle \chi_\downarrow | \Sigma_3 | \chi_\downarrow \rangle = +\frac{5}{3} \langle \chi_\uparrow | \Sigma_3 | \chi_\uparrow \rangle \quad (13)$$

$$g_T = \langle p \uparrow | \hat{u}^+ \beta \Sigma_3 \hat{u} - \hat{d}^+ \beta \Sigma_3 \hat{d} | p \uparrow \rangle = +\frac{4}{3} \langle \chi_\uparrow | \beta \Sigma_3 | \chi_\uparrow \rangle - \frac{1}{3} \langle \chi_\downarrow | \beta \Sigma_3 | \chi_\downarrow \rangle = +\frac{5}{3} \langle \chi_\uparrow | \beta \Sigma_3 | \chi_\uparrow \rangle \quad (14)$$

where we use obvious equalities:

$$\langle \chi_\uparrow | \Sigma_3 | \chi_\uparrow \rangle = -\langle \chi_\downarrow | \Sigma_3 | \chi_\downarrow \rangle, \quad \langle \chi_\uparrow | \beta \Sigma_3 | \chi_\uparrow \rangle = -\langle \chi_\downarrow | \beta \Sigma_3 | \chi_\downarrow \rangle \quad (15)$$

To proceed further, we choose the standard representation of the bispinor wave function

$$\Sigma_3 = \begin{pmatrix} \sigma_3 & 0 \\ 0 & \sigma_3 \end{pmatrix}, \quad \beta = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (16)$$

$$\chi_\uparrow(r, \theta, \phi) = \frac{1}{r} \begin{pmatrix} G(r) \Omega_{\frac{1}{2}, 0, \frac{1}{2}}(\theta, \phi) \\ F(r) \Omega_{\frac{1}{2}, 1, \frac{1}{2}}(\theta, \phi) \end{pmatrix} \quad (17)$$

$$\Omega_{\frac{1}{2}, 0, \frac{1}{2}}(\theta, \phi) = \begin{pmatrix} Y_{00}(\theta, \phi) \\ 0 \end{pmatrix}, \quad \Omega_{\frac{1}{2}, 1, \frac{1}{2}}(\theta, \phi) = \begin{pmatrix} -\sqrt{\frac{1}{3}} Y_{10}(\theta, \phi) \\ +\sqrt{\frac{2}{3}} Y_{11}(\theta, \phi) \end{pmatrix} \quad (18)$$

The following normalization conditions hold:

$$\int \Omega_{j'l'm'}^+ \Omega_{jlm} d\phi = \delta_{jj'} \delta_{ll'} \delta_{mm'}, \quad \int_0^\infty [G^2(r) + F^2(r)] dr = 1 \quad (19)$$

To take into account perturbative gluon exchanges we represent ΔH effectively as one-particle operators

$$\Delta H = \sum_{i=1}^3 \left(-\frac{\zeta}{r_i} \right) \quad (20)$$

³see [12] for similar calculations in bag models

Table 1: g_A , g_T , and η for various theoretical prescriptions in comparison with experimental data

	Exp.	NRQM	$\zeta = 0$	$\zeta = 0.3$
g_A	1.27	1.67	1.36	1.27
g_T	—	1.67	1.51	1.47
η	—	0	0.14	0.18

with an effective coupling constant ζ being fixed by realistic α_s values (see below), and the equations for $G(r)$, $F(r)$ acquire the form [13]

$$\begin{aligned} G' - \frac{1}{r}G - \left(\varepsilon + m + \sigma r + \frac{\zeta}{r} \right) F &= 0, \\ F' + \frac{1}{r}F + \left(\varepsilon - m - \sigma r + \frac{\zeta}{r} \right) G &= 0 \end{aligned} \quad (21)$$

where m is the light quark mass and ε is its energy.

Now, using equations (17,18), one can calculate explicitly the matrix elements appeared in (13,14):

$$\begin{aligned} \langle \chi_\uparrow | \Sigma_3 | \chi_\uparrow \rangle &= \\ &= \int r^2 dr d\Omega \left[\frac{G^2(r)}{r^2} \langle \Omega_{\frac{1}{2},0,\frac{1}{2}} | \sigma_3 | \Omega_{\frac{1}{2},0,\frac{1}{2}} \rangle + \frac{F^2(r)}{r^2} \langle \Omega_{\frac{1}{2},1,\frac{1}{2}} | \sigma_3 | \Omega_{\frac{1}{2},1,\frac{1}{2}} \rangle \right] = \\ &= \int dr \left(G^2(r) - \frac{1}{3} F^2(r) \right) \end{aligned} \quad (22)$$

$$\begin{aligned} \langle \chi_\uparrow | \beta \Sigma_3 | \chi_\uparrow \rangle &= \\ &= \int r^2 dr d\Omega \left[\frac{G^2(r)}{r^2} \langle \Omega_{\frac{1}{2},0,\frac{1}{2}} | \sigma_3 | \Omega_{\frac{1}{2},0,\frac{1}{2}} \rangle - \frac{F^2(r)}{r^2} \langle \Omega_{\frac{1}{2},1,\frac{1}{2}} | \sigma_3 | \Omega_{\frac{1}{2},1,\frac{1}{2}} \rangle \right] = \\ &= \int dr \left(G^2(r) + \frac{1}{3} F^2(r) \right) \end{aligned} \quad (23)$$

Finally g_A and g_T can be written as⁴

$$g_A = +\frac{5}{3} \left(1 - \frac{4}{3}\eta \right), \quad g_T = +\frac{5}{3} \left(1 - \frac{2}{3}\eta \right), \quad \eta = \int_0^\infty F^2(r) dr \quad (24)$$

Note, that in the nonrelativistic limit $\eta = 0$ and $g_A = g_T = 5/3$.

The light quark mass m is very small and can be omitted from (21) with good accuracy. So, the only dimensionful parameter σ remains in the model, but both g_A and g_T constants are dimensionless and, correspondingly, do not depend on σ . Thus, ζ proves to be the only essential model parameter. We have computed the values of η , g_A , and g_T for two

⁴It is clear from the aforesaid that neutron charges are simply opposite in sign to the proton ones, so we do not need to consider them separately

different values of ζ : $\zeta = 0$ and $\zeta = 0.3$. The results are given in Table 1; there also given for comparison the values obtained in the nonrelativistic quark model.

One can see that the resulting g_A is in the correct ballpark for $\zeta \in [0, 0.3]$. The value $\zeta = 0.3$ corresponds to the reasonable effective value of α_s in the qq potential, namely from $\left\langle \sum \frac{2}{3} \frac{\alpha_s}{r_{ij}} \right\rangle = \left\langle \sum \frac{\zeta}{r_i} \right\rangle$, and $\langle r_{ij} \rangle \approx \sqrt{3} \langle r_i \rangle$, one has

$$(\alpha_s)_{\text{eff}} = \frac{3\sqrt{3}}{4}\zeta \approx 0.39$$

and this value of α_s was checked in the actual calculation of the baryon masses in the Effective Hamiltonian approach [14]. It is rewarding that the resulting $g_A = 1.27$ is in close agreement with experiment.

Concerning the tensor charge g_T , one can compare these results with the lattice data [15], where both g_A and g_T are close to each other and are in the interval $1.12 \leq g_A, g_T \leq 1.18$ [15] for $m_\pi > 0.5$ GeV.

There are two possible unaccounted effects which can influence our results. First, the contribution of other terms in (10) – excited Dirac orbitals. The corresponding multichannel calculations done in [11] for magnetic moments, result in decreasing of the modulus of magnetic moments of proton and neutron by some 10 – 15% when one accounts for four Dirac orbitals for each quarks, and one can expect the same type of corrections for g_A . Second, the contribution of chiral degrees of freedom, i.e. of the π , η , K exchanges. Again, for nucleon magnetic moments these corrections are typically of the order of 10% [11], and we expect this to be an upper limit for g_A , since magnetic moments are much more sensitive to the contribution of the lowest Dirac components, than g_A and g_T , where these contributions enter quadratically and not linearly.

These corrections are not taken into account above, which is planned for a subsequent work, where also hyperon semileptonic decays are considered [16].

One should note, that relativistic approach to the baryon wave function, based on the light-cone formalism [17] was shown to improve the qualitative agreement of g_A with experiment, however quantitatively still far from experiment.

What is also possible to calculate in this approach are the baryon decuplet masses. We restrict ourselves here only to the ground state, so our leading S -wave approximation looks rather reasonable. As the spin and the isospin of any pair of light quarks in a decuplet baryon is equal to unity, one may expect the pion exchange effects are suppressed for decuplet masses. Indeed, calculations in [11] show that pion exchanges yield a negative shift to m_Δ by about 30 MeV. So, the baryon mass is completely determined by the eigenvalues of the system of equations (21). The only modification is due to the existence of a valence s -quark in a baryon – in fact, we should calculate independently the energy ε and orbital χ for the light quark q and for the strange quark s ; those are obtained from the same equations by an obvious replacement: $m \rightarrow m_i$, $\varepsilon \rightarrow \varepsilon_i$, $\chi \rightarrow \chi_i$, $i = q, s$.

However, on this way there still exists a problem for the full baryon wavefunction. We did not exclude clearly the baryon center-of-mass motion from the wavefunction and provided only the zero average value of the baryon 3-momentum, see (12). This means that the center-of-mass oscillates near zero point and bears a nonzero kinetic energy, which must be excluded from the baryon physical mass. The simplest way to do this is

to define the baryon mass in accordance with the Klein-Gordon equation:

$$M^2 = E^2 - \langle \mathbf{P}^2 \rangle, \quad (25)$$

where $\mathbf{P} = \sum \mathbf{p}_i$ is the total momentum of the three-quark system, and $E = \sum \varepsilon_i$ is the total energy.

As we consider baryons in the leading valence approximation with quarks in the S -wave orbitals, it is obvious that

$$\langle \mathbf{p}_i \mathbf{p}_j \rangle = 0 \text{ for } i \neq j, \text{ and } \langle \mathbf{p}_i^2 \rangle = \langle \chi_i | \mathbf{p}^2 | \chi_i \rangle \quad (26)$$

where χ_i the eigenfunction of the system of equations (21). Thus

$$\hat{H} \chi_i = \left(\boldsymbol{\alpha} \mathbf{p} + \beta \cdot m_i + \hat{1} \cdot \left(-\frac{\zeta}{r} \right) + \beta \cdot \sigma r \right) = \varepsilon_i \chi_i \quad (27)$$

and the averaged momentum squared can be expressed via two simple integrals:

$$\begin{aligned} \langle \chi_i | \mathbf{p}^2 | \chi_i \rangle &= \langle \chi_i | \boldsymbol{\alpha} \mathbf{p} | \boldsymbol{\alpha} \mathbf{p} | \chi_i \rangle = \left\langle \chi_i \left| \left(\varepsilon_i - \beta \cdot m_i - \hat{1} \cdot \left(-\frac{\zeta}{r} \right) - \beta \cdot \sigma r \right)^2 \right| \chi_i \right\rangle = \\ &= \left\langle \chi_i \left| \left(\varepsilon_i + \frac{\zeta}{r} \right)^2 + (m_i + \sigma r)^2 \right| \chi_i \right\rangle - 2 \left\langle \chi_i \left| \left(\varepsilon_i + \frac{\zeta}{r} \right) (m_i + \sigma r) \cdot \beta \right| \chi_i \right\rangle = \\ &= \int dr (G^2(r) + F^2(r)) \left(\left(\varepsilon_i + \frac{\zeta}{r} \right)^2 + (m_i + \sigma r)^2 \right) - \\ &\quad - 2 \int dr (G^2(r) - F^2(r)) \left(\varepsilon_i + \frac{\zeta}{r} \right) (m_i + \sigma r) \quad (28) \end{aligned}$$

Now the baryon mass can be calculated as follows:

$$M = \sqrt{\left(\sum \varepsilon_i \right)^2 - \sum \langle \mathbf{p}_i^2 \rangle} \quad (29)$$

As before, we neglect here the light quark mass, so, to make the numerical estimations, one needs to fix two dimensionful model parameters: the string tension σ and the strange quark mass m_s . Being guided by our previous calculations of baryon spectrum [14], we use the following values:

$$\sigma = 0.18 \text{ GeV}^2, \quad m_s = 210 \text{ MeV} \quad (30)$$

In the Table 2 we present the values of masses of ground-state baryons, calculated in this approach in comparison with experimental ones. One can see a fine agreement between the data.

This work was supported in part by the Grant for support of Leading Scientific Schools # 843.2006.2, by the State Contract # 02.445.11.7424, and in part by the RFBR grants ## 05-02-17869, 06-02-17120, 06-02-17012.

Table 2: Ground-state baryon masses, calculated in the Dirac orbital model, in comparison with the experimental values (in MeV)

Baryon	Δ	Σ^*	Ξ^*	Ω
Theor.	1233	1381	1527	1672
Exp.	1232	1383	1532	1672

References

- [1] T.D.Lee, *Particle Physics and Introduction to Field Theory* (Harwood Academic Publishers, 1981).
- [2] L.B.Okun, *Leptons and Quarks* (North Holland, 1982).
- [3] E.D.Commins and P.H.Bucksbaum, *Weak Interactions of Leptons and Quarks* (Cambridge University Press, 1983).
- [4] S.D.Bass, Eur. Phys. J. **A5**, 17 (1999).
- [5] S.Capstick, N.Isgur, Phys. Rev. D **34**, 2809 (1986).
- [6] U.Loering, K.Kretzschmar, B.C.Metsch, H.R.Petry, Eur. Phys. J. **A10**, 309 (2001).
- [7] V.Barone, A.Drago, P.C.Ratcliffe, Phys. Rep. **359**, 1 (2002).
- [8] L.Gamberg, G.Goldstein, Phys. Rev. Lett. **87**, 242001 (2001); hep-ph/0106178.
- [9] N.Cabibbo, E.C.Swallow, and R.Winston, hep-ph/0307214.
- [10] Yu.A.Simonov, Phys. Atom. Nucl. **62**, 1932 (1999).
- [11] Yu.A.Simonov, J.A.Tjon, J.Weda, Phys. Rev. D **65**, 094013 (2002); J.A.Tjon, J.Weda, Phys. Atom. Nucl. **68**, 591 (2005).
- [12] R.L.Jaffe and X.-D.Ji, Nucl. Phys. **B375**, 527 (1992).
- [13] V.D.Mur, V.S.Popov, Yu.A.Simonov, and V.P.Yurov, Sov. Phys – JETP **78**, 1 (1994).
- [14] I.M.Narodetskii and M.A.Trusov, Phys. Atom. Nucl. **67**, 762 (2004).
- [15] A.Ali-Khan et al. [QCDSF-UKQCD Collab.], Nucl. Phys. Proc. Suppl. **140**, 408 (2005); hep-lat/0409161.
- [16] Yu.A.Simonov and M.A.Trusov, in preparation.
- [17] F.Schlumpf, Phys. Rev. D **51**, 2262 (1995).